



UZEDY[®]
(risperidone) extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg
150 mg 200 mg 250 mg



**YOU'RE ALWAYS
THERE FOR THEM.
WE'RE HERE FOR YOU.**

Support, information, and resources to help you
as you care for someone living with schizophrenia.

FOR ADULTS LIVING WITH SCHIZOPHRENIA

People featured are not actual patients, caregivers, or doctors.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY[®]?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](#).

teva



YOUR ROLE AS A CAREGIVER MATTERS MORE THAN YOU KNOW

As a caregiver for an adult living with schizophrenia, your role is vital.

You might help your loved one manage medications, coordinate care appointments, monitor their symptoms, and offer essential emotional support.



PREVENTING RELAPSE MATTERS

Each episode can make symptoms harder to treat. After hospitalization, your role becomes even more vital. Consistent treatment reduces relapse risk and supports long-term stability.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not receive UZEDY® if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.uzedy.com).

A TREATMENT THAT MAY HELP PREVENT RELAPSES

Treatment for schizophrenia doesn't have to be a daily pill.
Long-acting injectables (LAIs) offer a different way.




UZEDY® IS THE FASTEST-GROWING BRANDED LAI

Based on the total volume of prescription claims of branded LAIs from May 2024 to May 2025.

WHAT MAKES UZEDY A DIFFERENT TYPE OF LAI?

UZEDY is delivered just beneath the skin. Most other schizophrenia LAIs are injected deep into muscle with a longer needle. UZEDY has one of the shortest LAI needles used to treat schizophrenia.*



This information should not be construed to imply differences in safety and efficacy between the products.

UZEDY SIGNIFICANTLY REDUCED THE RISK OF RELAPSE COMPARED WITH PLACEBO IN A CLINICAL STUDY.‡

Of the people who took once-monthly UZEDY, 7% (13 out of 183 people) had a relapse, while 29% of those who took a placebo (53 out of 181 people) had a relapse.



UZEDY is given every 1 or 2 months. It's absorbed rapidly after injection, within 24 hours. There are no starter doses or extra pills to remember to take.



In the clinical study, people taking UZEDY had a lower risk of relapse at 6 months than those who did not take UZEDY.§

*UZEDY is supplied as a prefilled syringe designed to provide 0.14 mL to 0.7 mL of medication, depending on the dose, with a 5/8-inch needle. No conclusions can be made between needle length and safety and effectiveness as this has not been studied.

†Shortest needle length; additional needle lengths for gluteal injection and/or patients with body weight over certain thresholds.

‡During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

§The relapse-free rate of people taking once-monthly UZEDY was 93% after 24 weeks (about 6 months) of treatment; 72% was the relapse-free rate of people taking a placebo.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including (continued):

- **Uncontrolled facial or body movements (tardive dyskinesia)** may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.uzedy.com).

All trademarks are the property of their respective owners.



CAREGIVERS NEED AND DESERVE SUPPORT, TOO

Being a caregiver for an adult living with schizophrenia can be a lot to handle. You might experience stress, frustration, or even burnout. It's important to remember to care for yourself as you care for your loved one.

ASK FOR HELP

See if someone can share responsibilities with you so you can carve out time for yourself.

- ✓ **How:** Make a list of tasks you could hand off, such as errands. Pick one and ask, "Would you be able to help with this sometime this week?"

SET BOUNDARIES

You may have many responsibilities besides caregiving, such as a career or other loved ones who rely on you. It's okay to say no and step back when you need space or support.

- ✓ **How:** Set clear times you're available. Turn off your phone during meals. Take 30 minutes for yourself, even just to sit in silence.

SHARE HOW YOU'RE FEELING

Consider talking to someone you trust—a friend, a relative, or a therapist—to share how you're feeling.

- ✓ **How:** Tell someone you trust, "I just need to talk for a few minutes." Or start with a quick message so you're not carrying it alone.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY® may cause serious side effects, including (continued):

- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.uzedy.com).

YOU'RE NOT ALONE. HELPFUL RESOURCES ARE AVAILABLE.

Connecting with others who understand your experience can be a great source of strength and insight. These organizations offer caregiver-specific resources, education, and support.

SUPPORT ORGANIZATIONS



NATIONAL ALLIANCE ON MENTAL ILLNESS (NAMI):

Provides education, advocacy, and support groups for caregivers.



Schizophrenia[™]
& Psychosis
Action Alliance

SCHIZOPHRENIA & PSYCHOSIS ACTION ALLIANCE:

Focused on better outcomes through education and caregiver empowerment.



CAREGIVER ACTION NETWORK:

Offers practical advice and community connection for those caring for loved ones with any health condition.

LEARN MORE ABOUT ALL YOUR SUPPORT OPTIONS

AND HOW TO SAVE ON UZEDY[®] AT [UZEDY.COM](https://uzedy.com)

ONGOING SUPPORT FROM TEVA

A patient support program is available to assist you and your loved one with planning for starting and continuing treatment with UZEDY by:

- [Confirming prescription coverage and pharmacy options based on eligibility and coverage]
- [Explaining Medicare and Medicaid coverage options]
- [Seeing what financial assistance options may be available]
- [Helping you understand prior authorization requirements]
- [Working with you to find an injection site near your home or work]
- [Providing over-the-phone nurse support to answer questions and help your loved one stay on UZEDY]

[Questions? Call [1-800-887-8100](tel:1-800-887-8100), Monday-Friday, 9 AM-8 PM ET.]

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including (continued):

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](https://uzedy.com).



APPROVED USE

UZEDY® (risperidone) extended-release injectable suspension is a prescription medicine used in adults for the treatment of schizophrenia.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Do not receive UZEDY if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.
- **Uncontrolled facial or body movements (tardive dyskinesia)** may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.
- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity,

family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.
- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**

Please see continued Important Safety Information on page 7 and the full Prescribing Information for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.uzedy.com).

- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.
- **Problems with control of your body temperature (too high or too low). Avoid getting overheated or dehydrated.**

The most common side effects of risperidone in patients with

- **Schizophrenia** included slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms.
- **Injection site reactions** including a lump or itching were reported with UZEDY®.

These are not all the possible side effects of UZEDY. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

Do not drink alcohol during treatment with UZEDY.

Before receiving UZEDY, tell your healthcare provider about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome.
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia).
- have diabetes or have a family history of diabetes.
- have had dizziness or fainting or are being treated for high blood pressure.

- have had a low white blood cell count.
- have or have had seizures or epilepsy.
- are pregnant or plan to become pregnant during treatment with UZEDY. It is not known if UZEDY will harm your unborn baby. Use of UZEDY during the third trimester of pregnancy may cause side effects in the newborn infant, including agitation, abnormal muscle tone, tremor, drowsiness, difficulty feeding, and difficulty breathing. Seek medical attention if you notice these signs. If you become pregnant during treatment with UZEDY, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics, or call 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. If you are receiving UZEDY and are breastfeeding, monitor your infant for sleepiness, inadequate weight gain, jitteriness, tremors, and abnormal muscle movements. Seek medical care if you notice these signs.
- have or have had kidney or liver problems.

Tell your healthcare provider about all the medicines you take or plan to take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. UZEDY and other medicines may affect each other.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For more information about UZEDY, see the full [Prescribing Information](#), including Boxed WARNING, or talk to your healthcare provider.


UZEDY[®]
(risperidone) extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg
150 mg 200 mg 250 mg

START THE CONVERSATION

Learning about the benefits of UZEDY[®] can help you support your loved one's journey with schizophrenia.

Talk to their doctor to see if UZEDY could be the right fit.



People featured are not actual patients, caregivers, or doctors.

Visit [UZEDY.com](https://www.uzedy.com) to learn more.

APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medicine used in adults for the treatment of schizophrenia.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information on previous pages and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.uzedy.com).

teva

© 2026 Teva Neuroscience, Inc.
RIS-41435 January 2026

YOUR ROLE AS A CAREGIVER MATTERS MORE THAN YOU KNOW

As a caregiver for an adult living with schizophrenia, your role is vital.

You might help your loved one manage medications, coordinate care appointments, monitor their symptoms, and offer essential emotional support.

PREVENTING RELAPSE MATTERS

Each episode can make symptoms harder to treat. After hospitalization, your role becomes even more vital. Consistent treatment reduces relapse risk and supports long-term stability.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Share how UZEDY® gives you a role in managing symptoms, or any of the components.

UZEDY® may cause serious side effects, including:

- Stroke in elderly people (contraindication problem) that can lead to death.
- Neuroleptic Malignant Syndrome (NMS) is a rare but very serious problem that can lead to death. Seek medical attention right away if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY®, including Boxed WARNING, at [UZEDY.com](#).

A TREATMENT THAT MAY HELP PREVENT RELAPSES

Treatment for schizophrenia doesn't have to be as difficult. Long-acting injectables (LAIs) offer a different way.

UZEDY® IS THE FASTEST-GROWING BRANDED LAI

What makes UZEDY a different choice for LAIs?

- UZEDY is delivered just once a month.
- LAIs are injected into the muscle with a longer needle. UZEDY has one of the shortest LAI needles available for schizophrenia.

What does this mean for you and your loved one?

- UZEDY is proven to reduce the risk of relapse compared with placebo in a clinical study.
- UZEDY is proven to be safe and effective for 12 months.
- UZEDY is proven to be safe and effective for 12 months.

UZEDY may cause serious side effects, including (continued):

- Stroke in elderly people (contraindication problem) that can lead to death.
- Neuroleptic Malignant Syndrome (NMS) is a rare but very serious problem that can lead to death. Seek medical attention right away if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY®, including Boxed WARNING, at [UZEDY.com](#).

CAREGIVERS NEED AND DESERVE SUPPORT, TOO

Being a caregiver for an adult living with schizophrenia can be a lot to handle. You might experience stress, frustration, or even burnout. It's important to remember to care for yourself as you care for your loved one.

ASK FOR HELP

Share how UZEDY® gives you a role in managing symptoms, or any of the components.

UZEDY® may cause serious side effects, including (continued):

- Stroke in elderly people (contraindication problem) that can lead to death.
- Neuroleptic Malignant Syndrome (NMS) is a rare but very serious problem that can lead to death. Seek medical attention right away if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY®, including Boxed WARNING, at [UZEDY.com](#).

YOU'RE NOT ALONE. HELPFUL RESOURCES ARE AVAILABLE.

Connecting with others who understand your experience can be a great source of strength and insight. These organizations offer caregiver-specific resources, education, and support.

SUPPORT ORGANIZATIONS

- NAMI** (National Alliance on Mental Illness) provides education, advocacy, and support for people with mental illness.
- SCHIZOPHRENIA & PSYCHOSES ACTION ALLIANCE** provides education, advocacy, and support for people with schizophrenia and psychosis.
- CAREGIVER ACTION NETWORK** offers practical advice and emotional support for caregivers.

LEARN MORE ABOUT ALL YOUR SUPPORT OPTIONS

AND HOW TO SAVE ON UZEDY® AT [UZEDY.com](#)

ONGOING SUPPORT FROM TEVA

A patient support program is available to assist you and your loved one with planning for starting and continuing treatment with UZEDY®.

- Confirming prescription coverage and plan many options based on eligibility and coverage.
- Exploring financial and medical assistance options.
- Seeking what financial assistance options may be available.
- Helping you understand your authorization requirements.
- Working with you to find an injection site near your home or work.
- Providing one-on-one phone support to answer questions and help your loved one get started on UZEDY®.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY® may cause serious side effects, including (continued):

- Stroke in elderly people (contraindication problem) that can lead to death.
- Neuroleptic Malignant Syndrome (NMS) is a rare but very serious problem that can lead to death. Seek medical attention right away if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY®, including Boxed WARNING, at [UZEDY.com](#).

APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medication used in adults for the treatment of schizophrenia.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY® may cause serious side effects, including an increased risk of death in elderly people who are frail, weak, or have other health problems. UZEDY® is not approved for use in patients with dementia-related psychosis.

Share how UZEDY® gives you a role in managing symptoms, or any of the components.

UZEDY® may cause serious side effects, including (continued):

- Stroke in elderly people (contraindication problem) that can lead to death.
- Neuroleptic Malignant Syndrome (NMS) is a rare but very serious problem that can lead to death. Seek medical attention right away if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY®, including Boxed WARNING, at [UZEDY.com](#).

APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medication used in adults for the treatment of schizophrenia.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY® may cause serious side effects, including an increased risk of death in elderly people who are frail, weak, or have other health problems. UZEDY® is not approved for use in patients with dementia-related psychosis.

Share how UZEDY® gives you a role in managing symptoms, or any of the components.

UZEDY® may cause serious side effects, including (continued):

- Stroke in elderly people (contraindication problem) that can lead to death.
- Neuroleptic Malignant Syndrome (NMS) is a rare but very serious problem that can lead to death. Seek medical attention right away if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY®, including Boxed WARNING, at [UZEDY.com](#).

START THE CONVERSATION

Learning about the benefits of UZEDY® can help you support your loved one's journey with schizophrenia.

Talk to your doctor to see if UZEDY® could be the right fit.

Visit [UZEDY.com](#) to learn more.

APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medication used in adults for the treatment of schizophrenia.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY® may cause serious side effects, including an increased risk of death in elderly people who are frail, weak, or have other health problems. UZEDY® is not approved for use in patients with dementia-related psychosis.

Share how UZEDY® gives you a role in managing symptoms, or any of the components.

UZEDY® may cause serious side effects, including (continued):

- Stroke in elderly people (contraindication problem) that can lead to death.
- Neuroleptic Malignant Syndrome (NMS) is a rare but very serious problem that can lead to death. Seek medical attention right away if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in blood pressure.

Please see additional Important Safety Information on previous pages and the full Prescribing Information for UZEDY®, including Boxed WARNING, at [UZEDY.com](#).