

**UZEDY**[®]
(risperidone)
extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg
150 mg 200 mg 250 mg

For adults with schizophrenia

DISCOVER WHAT FEWER SCHIZOPHRENIA RELAPSES* COULD DO FOR



U GOT THIS.

Help reduce your risk of relapse with **UZEDY**[®]

People featured are not actual patients, caregivers, or doctors.

*During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](#).





SCHIZOPHRENIA IS COMPLEX

UZEDY® MAY BE ABLE TO HELP REDUCE RELAPSES

UZEDY is a long-acting injectable (LAI) treatment for schizophrenia. With an LAI like UZEDY, you don't have to take a pill every day.



CONSISTENT TREATMENT MAY HELP PREVENT RELAPSES

When managing schizophrenia, it's common for people to forget to take their medication. This can cause relapses. Relapses can lower your chance of recovery and worsen your symptoms, so preventing them is important.*

*During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not receive UZEDY if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including (continued):

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](#).

HELP PREVENT RELAPSES WITH UZEDY^{®*}



LESS RISK OF RELAPSE

UZEDY significantly reduced the risk of relapse compared with placebo in a clinical study.*

Of the people who took once-monthly UZEDY, 7% (13 out of 183 people) had a relapse, while 29% of those who took a placebo (53 out of 181 people) had a relapse.



GREATER STABILITY

87% of people (159 out of 183) maintained stability[†] with their symptoms with once-monthly UZEDY vs 61% (110 out of 181) of people taking a placebo.



FEWER SETBACKS

The relapse-free rate of people taking once-monthly UZEDY was 93% after 24 weeks (about 6 months) of treatment; 72% was the relapse-free rate of people taking a placebo.

SAFETY WAS EXTENSIVELY STUDIED

The most common side effects of risperidone included slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms. Injection site reactions including a lump or itching were reported with UZEDY.

*During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

[†]In a clinical study of UZEDY, stability was defined based on patients who stayed outpatient, had few symptoms from a list of specific psychotic symptoms, and had symptom and clinical functioning scores below certain cutoffs for at least 4 weeks.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including (continued):

- **Uncontrolled facial or body movements (tardive dyskinesia)** may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at UZEDY.com.



UZEDY® IS AN LAI. BUT WHAT'S AN LAI?

An LAI is a treatment that releases medication slowly over a period of time. It comes in a shot, generally given by a doctor or nurse.

WITH UZEDY, YOU:

- ✓ Don't have to remember to take a pill every day
- ✓ Only get an injection once a month or every other month
- ✓ Don't need any additional starter doses or extra risperidone pills

92%

IN A SURVEY, 92% OF PEOPLE WHO TOOK UZEDY TO TREAT THEIR SCHIZOPHRENIA WERE SATISFIED. THEY RECEIVED UZEDY AS PART OF A TRIAL.*

*Data were collected from 63 patients, 24 physicians, and 25 nurses in a prospective, cross-sectional companion survey assessing the perceptions regarding ease of use and satisfaction with UZEDY. The survey was administered after a minimum of 2 experiences prescribing, administering, or receiving UZEDY.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including (continued):

- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](#).

UZEDY[®]
(risperidone)
extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg
150 mg 200 mg 250 mg

People featured are not actual patients, caregivers, or doctors.

UZEDY[®] HAS ONE OF THE SHORTEST LAI NEEDLES USED TO TREAT SCHIZOPHRENIA*

UZEDY is injected just below the skin, not deep into the muscle like other LAIs for schizophrenia. You can choose to get your UZEDY injection in your **upper arm or abdomen**.



UZEDY IS QUICKLY ABSORBED WITHIN 24 HOURS

You don't need any additional starter doses or risperidone pills once you begin UZEDY.



UZEDY KEEPS WORKING BETWEEN DOSES

UZEDY is slowly released over 1 or 2 months, until your next dose.



UZEDY HAS FLEXIBLE DOSING

You and your doctor can decide together if you'd rather get your UZEDY injection **every month or every other month**.

*UZEDY is supplied as a prefilled syringe designed to provide 0.14 mL to 0.7 mL of medication, depending on dose, with a 5/8-inch needle. No conclusions can be made between needle length and safety and effectiveness as this has not been studied.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including (continued):

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING**, at [UZEDY.com](#).



THINK UZEDY® MAY BE RIGHT FOR YOU?

Ask your doctor if they think so, too.

Here are some questions to get the conversation started.

- ? How do LAIs help people stay on track with their treatment?
- ? Am I a good candidate for an LAI?
- ? Is UZEDY a good option for me?
- ? How can UZEDY fit into my overall treatment plan?
- ? What are the possible benefits, risks, and side effects with UZEDY?
- ? If I were to start UZEDY, do you think I would get it once a month or every other month?
- ? Have your other patients tried UZEDY? What did they think of it?

Visit [UZEDY.com](https://www.uzedy.com) to download a full discussion guide to use at your next appointment.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including (continued):

- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.uzedy.com).



teva Total Support

- ✓ Helps you figure out what's covered by your insurance
- ✓ Shares financial assistance options and talks about what might be right for you
- ✓ Provides you with different tools and resources to keep you informed and confident at every step
- ✓ Helps you get in touch with your specialty pharmacy for smooth treatment delivery and personalized services
- ✓ Connects you with a nurse who will be there to support and guide you throughout your treatment

**YOU COULD SAVE ON UZEDY
ELIGIBLE PATIENTS
MAY PAY AS LITTLE
AS \$0 PER DOSE***



Visit [UZEDY.com](https://www.UZEDY.com) to learn more about your support options.

*Offer is available for patients with commercial insurance only. This offer is NOT available for patients eligible for Medicare, Medicaid, or any other form of government insurance coverage. Please see complete Terms and Conditions at www.UZEDY.com.

The Patient Savings Program for UZEDY (risperidone) extended-release injectable suspension is available to eligible patients who have been prescribed UZEDY and have commercial prescription insurance. Eligible patients may pay as little as \$0 for once-monthly or once-every-2-month dosing options of UZEDY. Maximum program assistance per prescription and annual benefit limits per individual apply and out-of-pocket expenses may vary. Patient is responsible for costs above maximum benefit amounts. This program is restricted to residents of the United States and United States territories. Uninsured and cash-paying patients are NOT eligible for this program. Patients enrolled in any state or federally funded healthcare program, including but not limited to, Medicare, Medigap, Medicaid, VA, DOD, TRICARE, Puerto Rico Government Health Insurance Plan, Medicare-eligible patients enrolled in an employer-sponsored health plan or prescription drug benefit program for retirees, are NOT eligible for this Program. Teva Neuroscience, Inc. and its affiliates reserve the right to change, rescind, revoke, or discontinue this program at any time without notice. Please see complete Terms and Conditions at www.UZEDY.com.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including (continued):

- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**
- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at [UZEDY.com](https://www.UZEDY.com).



APPROVED USE

UZEDY® (risperidone) extended-release injectable suspension is a prescription medicine used in adults for the treatment of schizophrenia.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Do not receive UZEDY if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.
- **Uncontrolled facial or body movements (tardive dyskinesia)** may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.
- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you

start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.
- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**
- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.

Please see continued Important Safety Information on page 9 and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING, at UZEDY.com.**



UZEDY® may cause serious side effects, including (continued):

- **Problems with control of your body temperature (too high or too low). Avoid getting overheated or dehydrated.**

The most common side effects of risperidone in patients with

- **Schizophrenia** included slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms.
- **Injection site reactions** including a lump or itching were reported with UZEDY.

These are not all the possible side effects of UZEDY. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

Do not drink alcohol during treatment with UZEDY.

Before receiving UZEDY, tell your healthcare provider about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome.
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia).
- have diabetes or have a family history of diabetes.
- have had dizziness or fainting or are being treated for high blood pressure.
- have had a low white blood cell count.
- have or have had seizures or epilepsy.

- are pregnant or plan to become pregnant during treatment with UZEDY. It is not known if UZEDY will harm your unborn baby. Use of UZEDY during the third trimester of pregnancy may cause side effects in the newborn infant, including agitation, abnormal muscle tone, tremor, drowsiness, difficulty feeding, and difficulty breathing. Seek medical attention if you notice these signs. If you become pregnant during treatment with UZEDY, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics, or call [1-866-961-2388](tel:1-866-961-2388) or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. If you are receiving UZEDY and are breastfeeding, monitor your infant for sleepiness, inadequate weight gain, jitteriness, tremors, and abnormal muscle movements. Seek medical care if you notice these signs.
- have or have had kidney or liver problems.

Tell your healthcare provider about all the medicines you take or plan to take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. UZEDY and other medicines may affect each other.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For more information about UZEDY, see the full [Prescribing Information](#), including **Boxed WARNING, or talk to your healthcare provider.**

UZEDY[®]
(risperidone) extended-release
injectable suspension

50 mg, 75 mg, 100 mg, 125 mg
150 mg, 200 mg, 250 mg

**FIND OUT WHAT
FEWER RELAPSES*
COULD DO FOR YOU.**

Talk to your doctor or visit
UZEDY.com to learn more.

People featured are not actual patients, caregivers, or doctors.

*During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY[®] once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

Visit UZEDY.com to learn more.

APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medicine used in adults for the treatment of schizophrenia.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information on previous pages and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING, at UZEDY.com.

teva

© 2026 Teva Neuroscience, Inc.
RIS-41805 January 2026